

# On trees with double domination number equal to total domination number plus one

Marcin Krzywkowski  
e-mail: marcin.krzywkowski@gmail.com

*Faculty of Applied Physics and Mathematics  
Gdańsk University of Technology  
Narutowicza 11/12, 80-233 Gdańsk, Poland*

## Abstract

A total dominating set of a graph  $G$  is a set  $D$  of vertices of  $G$  such that every vertex of  $G$  has a neighbor in  $D$ . A vertex of a graph is said to dominate itself and all of its neighbors. A double dominating set of a graph  $G$  is a set  $D$  of vertices of  $G$  such that every vertex of  $G$  is dominated by at least two vertices of  $D$ . The total (double, respectively) domination number of a graph  $G$  is the minimum cardinality of a total (double, respectively) dominating set of  $G$ . We characterize all trees with double domination number equal to total domination number plus one.

**Keywords:** total domination, double domination, tree.

**AMS Subject Classification:** 05C05, 05C69.

## 1 Introduction

Let  $G = (V, E)$  be a graph. By the neighborhood of a vertex  $v$  of  $G$  we mean the set  $N_G(v) = \{u \in V(G) : uv \in E(G)\}$ . The degree of a vertex  $v$ , denoted by  $d_G(v)$ , is the cardinality of its neighborhood. By a leaf we mean a vertex of degree one, while a support vertex is a vertex adjacent to a leaf. We say that a support vertex is strong (weak, respectively) if it is adjacent to at least two leaves (exactly one leaf, respectively). The path on  $n$  vertices we denote by  $P_n$ . By a star we mean a connected graph in which exactly one vertex has degree greater than one. By a double star we mean a graph obtained from a star by joining a positive number of vertices to one of its leaves. Let  $uv$  be an edge of a graph  $G$ . By subdividing the edge  $uv$  we mean removing it, and adding a new vertex, say  $x$ , along with two new edges  $ux$  and  $xv$ . Subdivided star is a graph obtained from a star by subdividing each one of its edges.

A subset  $D \subseteq V(G)$  is a dominating set of  $G$  if every vertex of  $V(G) \setminus D$  has a neighbor in  $D$ , while it is a total dominating set, abbreviated TDS, of  $G$  if every vertex of  $G$  has a neighbor in  $D$ . The domination (total domination, respectively) number of a graph  $G$ , denoted by  $\gamma(G)$  ( $\gamma_t(G)$ , respectively), is the minimum cardinality of a dominating (total dominating, respectively) set of  $G$ . Total domination in graphs was introduced by Cockayne, Dawes, and Hedetniemi [1]. For a comprehensive survey of domination in graphs, see [3, 4].

A vertex of a graph is said to dominate itself and all of its neighbors. A subset  $D \subseteq V(G)$  is a double dominating set, abbreviated DDS, of  $G$  if every vertex of  $G$  is dominated by at least two vertices of  $D$ . The double domination number of a graph  $G$ , denoted by  $\gamma_d(G)$ , is the minimum cardinality of a double dominating set of  $G$ . The study of double domination in graphs was initiated by Harary and Haynes [2].

A paired dominating set of a graph  $G$  is a dominating set of vertices whose induced subgraph has a perfect matching. The authors of [5] characterized all trees with equal total domination and paired domination numbers.

We characterize all trees with double domination number equal to total domination number plus one.

## 2 Results

Since the one-vertex graph does not have double dominating set, in this paper, by a tree we mean only a connected graph with no cycle, and which has at least two vertices.

We begin with the following four straightforward observations.

**Observation 1** *Every support vertex of a graph  $G$  is in every  $\gamma_t(G)$ -set.*

**Observation 2** *For every connected graph  $G$  of diameter at least three there exists a  $\gamma_t(G)$ -set that contains no leaf.*

**Observation 3** *Every leaf of a graph  $G$  is in every  $\gamma_d(G)$ -set.*

**Observation 4** *Every support vertex of a graph  $G$  is in every  $\gamma_d(G)$ -set.*

It is easy to see that  $\gamma_d(P_2) = \gamma_t(P_2) = 2$ . Now we prove that for every tree different than  $P_2$  the double domination number is greater than the total domination number.

**Lemma 5** *For every tree  $T \neq P_2$  we have  $\gamma_d(T) > \gamma_t(T)$ .*

**Proof.** Let  $n$  mean the number of vertices of the tree  $T$ . We proceed by induction on this number. Since  $T \neq P_2$ , we have  $\text{diam}(T) \geq 2$ . If  $\text{diam}(T) = 2$ , then  $T$  is a star  $K_{1,m}$ . We have  $\gamma_d(T) = m + 1 \geq 2 + 1 > 2 = \gamma_t(T)$ . Now let us assume that  $\text{diam}(T) = 3$ . Thus  $T$  is a double star. We have  $\gamma_d(T) = n \geq 4 > 2 = \gamma_t(T)$ .

Now assume that  $\text{diam}(T) \geq 4$ . Thus the order of the tree  $T$  is an integer  $n \geq 5$ . The result we obtain by the induction on the number  $n$ . Assume that the lemma is true for every tree  $T'$  of order  $n' < n$ .

First assume that some support vertex of  $T$ , say  $x$ , is strong. Let  $y$  and  $z$  mean leaves adjacent to  $x$ . Let  $T' = T - y$ . Let  $D'$  be any  $\gamma_t(T')$ -set. By Observation 1 we have  $x \in D'$ . Of course,  $D'$  is a TDS of the tree  $T$ . Thus  $\gamma_t(T) \leq \gamma_t(T')$ . Now let  $D$  be any  $\gamma_d(T)$ -set. By Observations 3 and 4 we have  $y, z, x \in D$ . It is easy to see that  $D \setminus \{y\}$  is a DDS of the tree  $T'$ . Therefore  $\gamma_d(T') \leq \gamma_d(T) - 1$ . Now we get  $\gamma_d(T) \geq \gamma_d(T') + 1 > \gamma_t(T') + 1 \geq \gamma_t(T) + 1 > \gamma_t(T)$ . Henceforth, we can assume that every support vertex of  $T$  is weak.

We now root  $T$  at a vertex  $r$  of maximum eccentricity  $\text{diam}(T)$ . Let  $t$  be a leaf at maximum distance from  $r$ ,  $v$  be the parent of  $t$ ,  $u$  be the parent of  $v$ , and  $w$  be the parent of  $u$  in the rooted tree. By  $T_x$  let us denote the subtree induced by a vertex  $x$  and its descendants in the rooted tree  $T$ .

First assume that  $d_T(u) \geq 3$ . Assume that  $u$  is adjacent to a leaf, say  $x$ . Let  $T' = T - T_v$ . Let  $D'$  be any  $\gamma_t(T')$ -set. By Observation 1 we have  $u \in D'$ . It is easy to see that  $D' \cup \{v\}$  is a TDS of the tree  $T$ . Thus  $\gamma_t(T) \leq \gamma_t(T') + 1$ . Now let  $D$  be any  $\gamma_d(T)$ -set. By Observations 3 and 4 we have  $t, x, v, u \in D$ . It is easy to see that  $D \setminus \{v, t\}$  is a DDS of the tree  $T'$ . Therefore  $\gamma_d(T') \leq \gamma_d(T) - 2$ . Now we get  $\gamma_d(T) \geq \gamma_d(T') + 2 > \gamma_t(T') + 2 \geq \gamma_t(T) + 1 > \gamma_t(T)$ .

Now assume that among the descendants of  $u$  there is a support vertex, say  $x$ , different than  $v$ . Let  $T' = T - T_v$ . Let  $D'$  be a  $\gamma_t(T')$ -set that contains no leaf. The vertex  $x$  has to have a neighbor in  $D'$ , thus  $u \in D'$ . It is easy to see that  $D' \cup \{v\}$  is a TDS of the tree  $T$ . Thus  $\gamma_t(T) \leq \gamma_t(T') + 1$ . Now let  $D$  be any  $\gamma_d(T)$ -set. By Observations 3 and 4 we have  $t, v, x \in D$ . If  $u \in D$ , then it is easy to see that  $D \setminus \{v, t\}$  is DDS of the tree  $T'$ . Now assume that  $u \notin D$ . Let us observe that  $D \cup \{u\} \setminus \{v, t\}$  is a DDS of the tree  $T'$ . Therefore  $\gamma_d(T') \leq \gamma_d(T) - 1$ . Now we get  $\gamma_d(T) \geq \gamma_d(T') + 1 > \gamma_t(T') + 1 \geq \gamma_t(T)$ .

Now assume that  $d_T(u) = 2$ . Let  $T' = T - T_u$ . If  $T' = P_2$ , then  $T = P_5$ . We have  $\gamma_d(P_5) = 4 > 3 = \gamma_t(P_5)$ . Now assume that  $T' \neq P_2$ . Let  $D'$  be any  $\gamma_t(T')$ -set. It is easy to see that  $D' \cup \{u, v\}$  is a TDS of the tree  $T$ . Thus  $\gamma_t(T) \leq \gamma_t(T') + 2$ . Now let us observe that there exists a  $\gamma_d(T)$ -set that does not contain the vertex  $u$ . Let  $D$  be such a set. By Observations 3 and 4 we have  $t, v \in D$ . Observe that  $D \setminus \{v, t\}$  is a DDS of the tree  $T'$ . Therefore  $\gamma_d(T') \leq \gamma_d(T) - 2$ . Now we get  $\gamma_d(T) \geq \gamma_d(T') + 2 > \gamma_t(T') + 2 \geq \gamma_t(T)$ . ■

Now we give a necessary condition for that the double domination number of a tree is equal to its total domination number plus one.

**Lemma 6** *If  $\gamma_d(T) = \gamma_t(T) + 1$ , then for every  $\gamma_d(T)$ -set  $D$ , every vertex of  $V(T) \setminus D$  has degree two.*

**Proof.** Suppose that there exists a  $\gamma_d(T)$ -set  $D$  that does not contain a vertex of  $T$ , say  $x$ , which has degree different than two. By Observation 3, every leaf belongs to the set  $D$ . Therefore  $d_T(x) \geq 3$ . First assume that some neighbor of  $x$ , say  $y$ , also does not belong to the set  $D$ . By  $T_1$  and  $T_2$  we denote the trees resulting from  $T$  by removing the edge  $xy$ . Let us observe that each one of those trees has at least three vertices. We define  $D_1 = D \cap V(T_1)$  and  $D_2 = D \cap V(T_2)$ . Let us observe that  $D_1$  is a DDS of the tree  $T_1$  and  $D_2$  is a DDS of the tree  $T_2$ . Let  $D'_1$  be any  $\gamma_t(T_1)$ -set and let  $D'_2$  be any  $\gamma_t(T_2)$ -set. By Lemma 5 we have  $\gamma_d(T_1) \geq \gamma_t(T_1) + 1$  and  $\gamma_d(T_2) \geq \gamma_t(T_2) + 1$ . Of course,  $D'_1 \cup D'_2$  is a TDS of the tree  $T$ . Thus  $\gamma_t(T) \leq |D'_1 \cup D'_2|$ . Now we get  $\gamma_d(T) = |D| = |D_1 \cup D_2| = |D_1| + |D_2| \geq \gamma_d(T_1) + \gamma_d(T_2) \geq \gamma_t(T_1) + 1 + \gamma_t(T_2) + 1 = |D'_1| + |D'_2| + 2 = |D'_1 \cup D'_2| + 2 \geq \gamma_t(T) + 2 > \gamma_t(T) + 1$ , a contradiction.

Now assume that all neighbors of  $x$  belong to the set  $D$ . First assume that there is a neighbor of  $x$ , say  $y$ , such that each one of the two trees resulting from  $T$  by removing the edge  $xy$  has at least three vertices. We get a contradiction similarly as when some neighbor of  $x$  does not belong to the set  $D$ . Now assume that there is no neighbor of  $x$  such that each one of the two trees resulting from  $T$  by removing the edge between them has at least three vertices. This implies that  $T$  is a subdivided star of order at least seven. Let  $n$  mean the number of vertices of the tree  $T$ . We have  $\gamma_d(T) = n - 1 = (n + 1)/2 + 1 + (n - 5)/2 = \gamma_t(T) + 1 + (n - 5)/2 > \gamma_t(T) + 1$ , a contradiction. ■

We characterize all trees with double domination number equal to total domination number plus one. For this purpose we introduce a family  $\mathcal{T} = \{P_3\} \cup \mathcal{A} \cup \mathcal{B}$ , where  $\mathcal{A} = \{A_1, A_2, \dots\}$  and  $\mathcal{B} = \{B_1, B_2, \dots\}$  are families of trees elements of which are given in Figure 1. A tree  $A_k$  has  $3k + 2$  vertices, and a tree  $B_k$  has  $3k + 3$  vertices.

Now we prove that for every tree of the family  $\mathcal{T}$ , the double domination number is equal to the total domination number plus one.

**Lemma 7** *If  $T \in \mathcal{T}$ , then  $\gamma_d(T) = \gamma_t(T) + 1$ .*

**Proof.** Of course,  $\gamma_d(P_3) = 3 = 2 + 1 = \gamma_t(P_3) + 1$ . Let  $k$  be a positive integer. For trees  $A_k$  and  $B_k$  we consider the labeling of the vertices as in Figure 1.

Let  $D$  be a  $\gamma_t(A_k)$ -set that contains no leaf. By Observation 1 we have




Figure 1: The path  $P_3$ , a tree  $A_k$  of the family  $\mathcal{A}$ , and a tree  $B_k$  of the family  $\mathcal{B}$

$b_1, b_2, \dots, b_k, x \in D$ . Since each one of the vertices  $b_1, b_2, \dots, b_k$  has to have a neighbor in the set  $D$ , we have  $a_1, a_2, \dots, a_k \in D$ . Therefore  $\gamma_t(A_k) \geq 2k + 1$ . It is easy to observe that  $\{b_1, c_1, b_2, c_2, \dots, b_k, c_k, x, y\}$  is a DDS of the tree  $A_k$ . Thus  $\gamma_d(A_k) \leq 2k + 2$ . Now we get  $\gamma_d(A_k) \leq 2k + 2 \leq \gamma_t(A_k) + 1$ . On the other hand, by Lemma 5 we have  $\gamma_d(A_k) \geq \gamma_t(A_k) + 1$ .

Now let  $D$  be a  $\gamma_t(B_k)$ -set that contains no leaf. By Observation 1 we have  $b_1, b_2, \dots, b_k, y \in D$ . Since each one of the vertices  $b_1, b_2, \dots, b_k, y$  has to have a neighbor in  $D$ , we have  $a_1, a_2, \dots, a_k, x \in D$ . Therefore  $\gamma_t(B_k) \geq 2k + 2$ . It is easy to observe that  $\{b_1, c_1, b_2, c_2, \dots, b_k, c_k, x, y, z\}$  is a DDS of the tree  $B_k$ . Thus  $\gamma_d(B_k) \leq 2k + 3$ . Now we get  $\gamma_d(B_k) \leq 2k + 3 \leq \gamma_t(B_k) + 1$ . This implies that  $\gamma_d(B_k) = \gamma_t(B_k) + 1$ . ■

Now we prove that if the double domination number of a tree is equal to its total domination number plus one, then the tree belongs to the family  $\mathcal{T}$ .

**Lemma 8** *Let  $T$  be a tree. If  $\gamma_d(T) = \gamma_t(T) + 1$ , then  $T \in \mathcal{T}$ .*

**Proof.** Let  $n$  mean the number of vertices of the tree  $T$ . We proceed by induction on this number. If  $\text{diam}(T) = 1$ , then  $T = P_2$ . We have  $\gamma_d(T) = 2 = \gamma_t(T) \neq \gamma_t(T) + 1$ . If  $\text{diam}(T) = 2$ , then  $T$  is a star  $K_{1,m}$ . If  $T = P_3$ , then  $T \in \mathcal{T}$ . Now assume that  $T$  is a star different than  $P_3$ . We have  $\gamma_d(T) = m + 1 \geq 3 + 1 > 2 + 1 = \gamma_t(T) + 1$ . Now let us assume that  $\text{diam}(T) = 3$ . Thus  $T$  is a double star. We have  $\gamma_d(T) = n \geq 4 > 3 = 2 + 1 = \gamma_t(T) + 1$ .

Now assume that  $\text{diam}(T) \geq 4$ . Thus the order of the tree  $T$  is an integer  $n \geq 5$ . The result we obtain by the induction on the number  $n$ . Assume that the lemma is true for every tree  $T'$  of order  $n' < n$ .

First assume that some support vertex of  $T$ , say  $x$ , is strong. Let  $y$  and  $z$  mean leaves adjacent to  $x$ . Let  $T' = T - y$ . Let  $D'$  be any  $\gamma_t(T')$ -set. By Observation 1 we have  $x \in D'$ . Of course,  $D'$  is a TDS of the tree  $T$ . Thus  $\gamma_t(T) \leq \gamma_t(T')$ . Now let  $D$  be any  $\gamma_d(T)$ -set. By Observations 3 and 4 we have  $y, z, x \in D$ . It is easy to see that  $D \setminus \{y\}$  is a DDS of the tree  $T'$ . Therefore  $\gamma_d(T') \leq \gamma_d(T) - 1$ . Now we get  $\gamma_d(T') \leq \gamma_d(T) - 1 = \gamma_t(T) \leq \gamma_t(T')$ . This is a contradiction as by Lemma 5 we have  $\gamma_d(T') > \gamma_t(T')$ . Thus every support vertex of  $T$  is weak.

We now root  $T$  at a vertex  $r$  of maximum eccentricity  $\text{diam}(T)$ . Let  $t$  be a leaf at maximum distance from  $r$ ,  $v$  be the parent of  $t$ ,  $u$  be the parent of  $v$ , and  $w$  be the parent of  $u$  in the rooted tree. By  $T_x$  let us denote the subtree induced by a vertex  $x$  and its descendants in the rooted tree  $T$ .

First assume that  $d_T(u) \geq 3$ . Assume that  $u$  is adjacent to a leaf, say  $x$ . Let  $T' = T - T_v$ . Let  $D'$  be any  $\gamma_t(T')$ -set. By Observation 1 we have  $u \in D'$ . It is easy to see that  $D' \cup \{v\}$  is a TDS of the tree  $T$ . Thus  $\gamma_t(T) \leq \gamma_t(T') + 1$ . Now let  $D$  be any  $\gamma_d(T)$ -set. By Observations 3 and 4 we have  $t, x, v, u \in D$ . It is easy to see that  $D \setminus \{v, t\}$  is a DDS of the tree  $T'$ . Therefore  $\gamma_d(T') \leq \gamma_d(T) - 2$ . Now we get  $\gamma_d(T') \leq \gamma_d(T) - 2 = \gamma_t(T) - 1 \leq \gamma_t(T')$ , a contradiction.

Thus every descendant of  $u$  is a support vertex. Let  $x$  mean a child of  $u$  different than  $v$ . Let  $T' = T - T_v$ . Let  $D'$  be a  $\gamma_t(T')$ -set that contains no leaf. The vertex  $x$  has to have a neighbor in  $D'$ , thus  $u \in D'$ . It is easy to see that  $D' \cup \{v\}$  is a TDS of the tree  $T$ . Thus  $\gamma_t(T) \leq \gamma_t(T') + 1$ . Now let  $D$  be any  $\gamma_d(T)$ -set. By Observations 3 and 4 we have  $t, v, x \in D$ . By Lemma 6 we have  $u \in D$ . It is easy to see that  $D \setminus \{v, t\}$  is a DDS of the tree  $T'$ . Therefore  $\gamma_d(T') \leq \gamma_d(T) - 2$ . Now we get  $\gamma_d(T') \leq \gamma_d(T) - 2 = \gamma_t(T) - 1 \leq \gamma_t(T')$ , a contradiction.

Now assume that  $d_T(u) = 2$ . Let  $T' = T - T_u$ . If  $T' = P_2$ , then  $T = P_5$ . Obviously,  $P_5 = A_1 \in \mathcal{T}$ . Now assume that  $T' \neq P_2$ . Let  $D'$  be any  $\gamma_t(T')$ -set. It is easy to see that  $D' \cup \{u, v\}$  is a TDS of the tree  $T$ . Thus  $\gamma_t(T) \leq \gamma_t(T') + 2$ . Now let us observe that there exists a  $\gamma_d(T)$ -set that does not contain the vertex  $u$ . Let  $D$  be such a set. By Observations 3 and 4 we have  $t, v \in D$ . Observe that  $D \setminus \{v, t\}$  is a DDS of the tree  $T'$ . Therefore  $\gamma_d(T') \leq \gamma_d(T) - 2$ . Now we get  $\gamma_d(T') \leq \gamma_d(T) - 2 = \gamma_t(T) - 1 \leq \gamma_t(T') + 1$ . This implies that  $\gamma_d(T') = \gamma_t(T') + 1$ . By the inductive hypothesis we have  $T' \in \mathcal{T}$ . If  $T' = P_3$ , then  $T = P_6$ . Obviously,  $P_6 = B_1 \in \mathcal{T}$ . Now assume that  $T' \neq P_3$ . We distinguish between the following two cases:  $T' \in \mathcal{A}$  and  $T' \in \mathcal{B}$ .

**Case 1.**  $T' \in \mathcal{A}$ . Let  $T' = A_k$ . We consider the labeling of the vertices as in Figure 1. If  $w$  corresponds to  $x$ , then it is easy to observe that  $T = A_{k+1} \in \mathcal{T}$ .

Now assume that  $w$  corresponds to  $y$ . It is easy to see that  $\{a_1, b_1, a_2, b_2, \dots, a_k, b_k, u, v\}$  is a TDS of the tree  $T$ . Thus  $\gamma_t(T) \leq 2k + 2$ . Now let

$D$  be any  $\gamma_d(T)$ -set. By Observations 3 and 4 we have  $c_1, b_1, c_2, b_2, \dots, c_k, b_k, t, v \in D$ . By Lemma 6 we have  $x \in D$ . It is easy to see that those vertices do not form a DDS of the tree  $T$ . Therefore  $\gamma_d(T) \geq 2k + 4$ . Now we get  $\gamma_d(T) \geq 2k + 4 > 2k + 3 \geq \gamma_t(T) + 1$ , a contradiction.

Now assume that  $w$  corresponds to  $a_i$ , for some  $i$ . It is easy to see that  $\{a_1, b_1, a_2, b_2, \dots, a_k, b_k, x, u, v\}$  is a TDS of the tree  $T$ . Thus  $\gamma_t(T) \leq 2k + 3$ . Now let  $D$  be any  $\gamma_d(T)$ -set. By Observations 3 and 4 we have  $c_1, b_1, c_2, b_2, \dots, c_k, b_k, y, x, t, v \in D$ . By Lemma 6 we have  $a_i \in D$ . Therefore  $\gamma_d(T) \geq 2k + 5$ . Now we get  $\gamma_d(T) \geq 2k + 5 > 2k + 4 \geq \gamma_t(T) + 1$ , a contradiction.

Now assume that  $w$  corresponds to  $b_i$ , for some  $i$ . Let us observe that  $\{a_1, b_1, a_2, b_2, \dots, a_{i-1}, b_{i-1}, b_i, a_{i+1}, b_{i+1}, \dots, a_k, b_k, x, u, v\}$  is a TDS of the tree  $T$ . Thus  $\gamma_t(T) \leq 2k + 2$ . Now let  $D$  be any  $\gamma_d(T)$ -set. By Observations 3 and 4 we have  $c_1, b_1, c_2, b_2, \dots, c_k, b_k, y, x, t, v \in D$ . Therefore  $\gamma_d(T) \geq 2k + 4$ . Now we get  $\gamma_d(T) \geq 2k + 4 > 2k + 3 \geq \gamma_t(T) + 1$ , a contradiction.

Now assume that  $w$  corresponds to  $c_i$ , for some  $i$ . Observe that  $\{a_1, b_1, a_2, b_2, \dots, a_{i-1}, b_{i-1}, a_i, a_{i+1}, b_{i+1}, \dots, a_k, b_k, x, u, v\}$  is a TDS of the tree  $T$ . Thus  $\gamma_t(T) \leq 2k + 2$ . Now let  $D$  be any  $\gamma_d(T)$ -set. By Observations 3 and 4 we have  $c_1, b_1, c_2, b_2, \dots, c_{i-1}, b_{i-1}, c_{i+1}, b_{i+1}, \dots, c_k, b_k, y, x, t, v \in D$ . Observe that adding any one of the remaining vertices to those vertices does not give us a DDS of the tree  $T$ . Therefore  $\gamma_d(T) \geq 2k + 4$ . Now we get  $\gamma_d(T) \geq 2k + 4 > 2k + 3 \geq \gamma_t(T) + 1$ , a contradiction.

**Case 2.**  $T' \in \mathcal{B}$ . Let  $T' = B_k$ . Let us consider the labeling of the vertices as in Figure 1. If  $w$  corresponds to  $x$ , then it is easy to see that  $T = B_{k+1} \in \mathcal{T}$ .

Now assume that  $w$  corresponds to  $z$ . Observe that  $\{a_1, b_1, a_2, b_2, \dots, a_k, b_k, z, u, v\}$  is a TDS of the tree  $T$ . Thus  $\gamma_t(T) \leq 2k + 3$ . Now let  $D$  be any  $\gamma_d(T)$ -set. By Observations 3 and 4 we have  $c_1, b_1, c_2, b_2, \dots, c_k, b_k, t, v \in D$ . By Lemma 6 we have  $x \in D$ . Let us observe that adding any one of the remaining vertices to those vertices does not give us a DDS of the tree  $T$ . Therefore  $\gamma_d(T) \geq 2k + 5$ . Now we get  $\gamma_d(T) \geq 2k + 5 > 2k + 4 \geq \gamma_t(T) + 1$ , a contradiction.

Now assume that  $w$  corresponds to  $y$ . Observe that  $\{a_1, b_1, a_2, b_2, \dots, a_k, b_k, y, u, v\}$  is a TDS of the tree  $T$ . Thus  $\gamma_t(T) \leq 2k + 3$ . Now let  $D$  be any  $\gamma_d(T)$ -set. By Observations 3 and 4 we have  $c_1, b_1, c_2, b_2, \dots, c_k, b_k, z, y, t, v \in D$ . By Lemma 6 we have  $x \in D$ . Therefore  $\gamma_d(T) \geq 2k + 5$ . Now we get  $\gamma_d(T) \geq 2k + 5 > 2k + 4 \geq \gamma_t(T) + 1$ , a contradiction.

Now assume that  $w$  corresponds to  $a_i$ , for some  $i$ . Observe that  $\{a_1, b_1, a_2, b_2, \dots, a_k, b_k, x, y, u, v\}$  is a TDS of the tree  $T$ . Thus  $\gamma_t(T) \leq 2k + 4$ . Now let  $D$  be any  $\gamma_d(T)$ -set. By Observations 3 and 4 we have  $c_1, b_1, c_2, b_2, \dots, c_k, b_k, z, y, t, v \in D$ . By Lemma 6 we have  $x, a_i \in D$ . Therefore  $\gamma_d(T) \geq 2k + 6$ . Now we get  $\gamma_d(T) \geq 2k + 6 > 2k + 5 \geq \gamma_t(T) + 1$ , a contradiction.

Now assume that  $w$  corresponds to  $b_i$ , for some  $i$ . Let us observe that  $\{a_1, b_1, a_2, b_2, \dots, a_{i-1}, b_{i-1}, b_i, a_{i+1}, b_{i+1}, \dots, a_k, b_k, x, y, u, v\}$  is a TDS of the tree  $T$ . Thus  $\gamma_t(T) \leq 2k + 3$ . Now let  $D$  be any  $\gamma_d(T)$ -set. By Observations 3 and 4 we have  $c_1, b_1, c_2, b_2, \dots, c_k, b_k, z, y, t, v \in D$ . By Lemma 6 we have  $x \in D$ . Therefore  $\gamma_d(T) \geq 2k + 5$ . Now we get  $\gamma_d(T) \geq 2k + 5 > 2k + 4 \geq \gamma_t(T) + 1$ , a contradiction.

Now assume that  $w$  corresponds to  $c_i$ , for some  $i$ . Let us observe that  $\{a_1, b_1, a_2, b_2, \dots, a_{i-1}, b_{i-1}, a_i, a_{i+1}, b_{i+1}, \dots, a_k, b_k, x, y, u, v\}$  is a TDS of the tree  $T$ . Thus  $\gamma_t(T) \leq 2k + 3$ . Now let  $D$  be any  $\gamma_d(T)$ -set. By Observations 3 and 4 we have  $c_1, b_1, c_2, b_2, \dots, c_{i-1}, b_{i-1}, c_{i+1}, b_{i+1}, \dots, c_k, b_k, z, y, t, v \in D$ . By Lemma 6 we have  $x \in D$ . Observe that adding any one of the remaining vertices to those vertices does not give us a DDS of the tree  $T$ . Therefore  $\gamma_d(T) \geq 2k + 5$ . Now we get  $\gamma_d(T) \geq 2k + 5 > 2k + 4 \geq \gamma_t(T) + 1$ , a contradiction. ■

As an immediate consequence of Lemmas 7 and 8, we have the following characterization of the trees with double domination number equal to total domination number plus one.

**Theorem 9** *Let  $T$  be a tree. Then  $\gamma_d(T) = \gamma_t(T) + 1$  if and only if  $T \in \mathcal{T}$ .*

## References

- [1] E. Cockayne, R. Dawes, and S. Hedetniemi, *Total domination in graphs*, Networks 10 (1980), 211–219.
- [2] F. Harary, T. Haynes, *Double domination in graphs*, Ars Combinatoria 55 (2000), 201–213.
- [3] T. Haynes, S. Hedetniemi, and P. Slater, *Fundamentals of Domination in Graphs*, Marcel Dekker, New York, 1998.
- [4] T. Haynes, S. Hedetniemi, and P. Slater (eds.), *Domination in Graphs: Advanced Topics*, Marcel Dekker, New York, 1998.
- [5] M. Henning, L. Kang, and E. Shan, *A characterization of trees with equal total domination and paired-domination numbers*, Australasian Journal of Combinatorics 30 (2004), 31–39.